

Enter Document Control Number or
Bar Code

60070002186

United States Environmental Protection Agency
Washington, DC 20460


ORIGINAL

Document Description

Focus Notes L07-0295

Date

7-19-07



Draft

FOCUS REPORT
New Chemicals Program

PART I: BACKGROUND

Written By: KMB

FOCUS DATE: 7/9/2007

FOCUS CHAIR: D. Jones

COMPANY: Tracerco

CASE NUMBER(S): L07-0295 through and

PART II: SAT RESULTS

HEALTH: 1-2

ECOTOX: 1

OCCUPATIONAL
EXPOSURE: 2-3A

CONSUMER
EXPOSURE: -

ENVIRONMENTAL
RELEASES: -

Additional SAT
Information:

PART III: OTHER FACTORS

- a. PRODUCTION VOLUME: 500 kg/yr
- b. PROD VOL OTHER: ***Binding Option Marked***
LVE Assessed at Production Volume
- c. USE: Tracer chemical to measure flow in deep oil-bearing strata
- d. REGULATORY HISTORY: L07-0271 (F) ☐ GRANTED WITH CONDITION

e. TEST DATA:

f. IMPORTED ☒

MANUFACTURED

BOTH ☐

g. MSDS: ☒

h. CATEGORY: Neutral Organics

CATEGORY 2:

PART IV: SUMMARY OF SAT ASSESSMENT

CASE NUMBERS: L07-0290 to L07-0330

FATE: MW162 to 216

solids with mp for L070271 (F) = 126 °C (M)

log Kow for the free acid = 1.8 to 2.9 (ClogP), 1.2 to 2.8 (EPI), 1.59 to 3.10 (M)

log Kow for L070271 = -5.87 with pH? (HPLC)

log Kow for L070296 = -2.11 with pH? (HPLC)

log Kow for L070297 = -3.62 with pH? (HPLC)

log Kow for L070298 = -1.64 with pH? (HPLC)

S > 200 mg/L to > 10 g/L at 20 °C (P)

vp < 1.0E-6 mm Hg or torr at 25 °C (P)

bp = 460 °C (P)

H for the covalent ion pair = 1.2E-7 to 9.5E-7 (P)

log Koc for the covalent ion pair = 1.4 to 2.0 (P)

log fish BCF = 0.50 (P)



sorption to sludge = low (P)

test data for L070271 for aerobic biodegradation in seawater at 20 C, via closed bottle (OECD306) were:

time biodegradation
(d) (percent)

5 0
14 2
28 63

test data for aerobic biodegradation for the [REDACTED] of L070299 from [REDACTED] were:
15% biodegradation in 28 d, thus, not readily biodegradable via CO2 evolution in modified Sturm test (OECD301B); if test result is due solely to ester hydrolysis and degradation of the [REDACTED], then removal via POTW of the parent would be >= 90% but notifier did not measure degradation products;

POTW removal = 0% to 90 via sorption and possible biodegradation
time for complete ultimate aerobic biodegradation = weeks to => months
sorption to soils and sediments = low (P)
PBT Potential: P2B1T2 to P3B1T2

HEALTH: Absorption nil thru skin based on physical/chemical properties; good thru lungs based on analogs; and good thru the GI tract based on analogs;

test data for the methyl ester of L070299, [REDACTED], were:
rat acute oral LD50 = 800 mg/kg with toxic signs; LD100 = 2 g/kg, LD0 = 300 mg/kg;
rat acute dermal LD0 = 2.0 g/kg with no toxic signs;
slight and transient (2 d) skin irritation in rabbits;
slight and transient (1 d) eye irritation in rabbits;
Ames test was negative;
E. coli test was negative;
chromosome aberration test with V79 cells was positive with activation, but negative without activation;
no skin sensitization in guinea pigs (M&K);
rat 28-d subchronic oral-gavage with doses = 1000, 300, and 100 mg/kg/d with NOAEL = 100 mg/kg/d and LOEL = 300 mg/kg/d based on salivation and increased water consumption; effects at 1000 mg/kg/d were slight to severe salivation, unsteady gait, motor activity significantly decreased and effects to the liver and kidneys;

concern for asthma and developmental toxicity based on data for benzoic acid, note: the mechanism for the asthma is unknown;

concern for possible mutagenicity, liver toxicity, and kidney toxicity based on data for [REDACTED] which was the [REDACTED] of L070299, however, the [REDACTED] will have some acylating activity which is absent in the acid, thus, the acid will be less toxic than the [REDACTED];

low to moderate concern for toxicity

ECOTOX: Predicted (P) and measured (M) toxicity values in mg/L (ppm) are:

fish 96-h LC50	> 100.0	P
SW fish 96-h LC50	= 440.0	M S,N L070271
SW fish 96-h LC50	> 320.0	M S,N L070290
SW fish 96-h LC50	> 320.0	M S,N L070291
daphnid 48-h LC50	> 100.0	P
SW invert Ac ton 48-h LC50	= 2830.0	M S,N L070271
SW invert Ac ton 48-h LC50	= 1500.0	M S,N L070290
SW invert Ac ton 48-h LC50	= 430.0	M S,N L070291

SW invert Ac ton 48-h LC50 = 480.0 M S,N L070292
 SW invert Ac ton 48-h LC50 = 270.0 M S,N L070293
 SW invert Ac ton 48-h LC50 = 250.0 M S,N L070294
 SW invert Ac ton 48-h LC50 = 250.0 M S,N L070295
 SW invert Ac ton 48-h LC50 = 300.0 M S,N L070296
 SW invert Ac ton 48-h LC50 = 430.0 M S,N L070297
 SW invert Ac ton 48-h LC50 = 440.0 M S,N L070298
 SW invert Ac ton 48-h LC50 = 170.0 M S,N L070299
 SW invert Ac ton 48-h LC50 = 130.0 M S,N L070300
 green algal 96-h EC50 > 100.0 P
 SW algae Sk cost 72-h EC50 c = 250.0 M S,N L070271
 SW algae Sk cost 72-h EC50 r > 10000.0 M S,N L070290
 SW algae Sk cost 72-h EC50 r = 430.0 M S,N L070291
 SW algae Sk cost 72-h EC50 r = 660.0 M S,N L070292
 SW algae Sk cost 72-h EC50 r = 2100.0 M S,N L070296
 SW algae Sk cost 72-h EC50 r = 1500.0 M S,N L070297
 SW algae Sk cost 72-h EC50 r = 700.0 M S,N L070300
 fish chronic value > 10.0 P
 daphnid ChV > 10.0 P
 algal ChV > 10.0 P
 SW algae Sk cost ChV c = 100.0 M S,N L070271
 SW algae Sk cost ChV r = 5600.0 M S,N L070290
 SW algae Sk cost ChV r < 100.0 M S,N L070291
 SW algae Sk cost ChV r = 320.0 M S,N L070292
 SW algae Sk cost ChV r = 1000.0 M S,N L070296
 SW algae Sk cost ChV r = 320.0 M S,N L070297
 SW algae Sk cost ChV r = 320.0 M S,N L070300

benthic

SW invert Coror vol 10-d LC50 = 6558.0 mg/kg DWT M S,N L070271
 SW invert Coror vol 10-d NOEC = 470.0 mg/kg DWT M S,N L070271

SW invert Coror vol 10-d LC50 = 7300.0 mg/kg DWT M S,N L070290
 SW invert Coror vol 10-d NOEC = 1400.0 mg/kg DWT M S,N L070290

SW invert Coror vol 10-d LC50 = 3800.0 mg/kg DWT M S,N L070291
 SW invert Coror vol 10-d NOEC = 150.0 mg/kg DWT M S,N L070291

SW invert Coror vol 10-d LC50 = 6700.0 mg/kg DWT M S,N L070292
 SW invert Coror vol 10-d NOEC = 1400.0 mg/kg DWT M S,N L070292

SW invert Coror vol 10-d LC50 = 410.0 mg/kg DWT M S,N L070296
 SW invert Coror vol 10-d NOEC = 130.0 mg/kg DWT M S,N L070296

SW invert Coror vol 10-d LC50 = 330.0 mg/kg DWT M S,N L070297
 SW invert Coror vol 10-d NOEC = 160.0 mg/kg DWT M S,N L070297

SW invert Coror vol 10-d LC50 = 280.0 mg/kg DWT M S,N L070300
 SW invert Coror vol 10-d NOEC = 16.0 mg/kg DWT M S,N L070300

Predictions are based on SARs for neutral organic chemicals with 10X less toxicity due to the substitution of the acid, or SARs for anionic surfactants-carboxylic acid-C4.Na; SAR chemical class = surfactant-anionic-F1 to F4 and CF3 benzene-COO.Na; MW162 to 216; solids with mp for L070271 (F) = 126 °C (M); log Kow for the free acid = 1.8 to 2.9 (ClogP), 1.2 to 2.8 (EPI), 1.59 to 3.10 (M); log Kow for L070271 = -5.87 with pH? (HPLC); S > 200 mg/L at 20 °C (P); pH7; effective concentrations based on 100% active ingredients and mean measured concentrations; hardness <150.0 mg/L as CaCO3; and TOC <2.0 mg/L;
 low concern for toxicity
 assessment factor = 10.0
 concern concentration = 1.0 mg/L (ppm)

P2REC: forward to FOCUS with support.

PART V: RAD RISK RATIONALE: HUMAN HEALTH

PART VI: SUMMARY OF EXPOSURE/RELEASE

Proc:

1 site, 3 workers, 10 d/yr

Inhalation: Particulate: 1.4×10^2 mg/d

Dermal: Not Required per SAT

Releases to Water 1: 1.0 kg/s/d, 10 d/yr

Releases to Water 2: 5.0×10^{-1} kg/s/d, 10 d/yr

OR Air OR Incineration OR Landfill

Releases to Water 3: 1.0 kg/s/d, 10 d/yr

OR Incineration OR Landfill

Fate: Releases to Water 1, 2, & 3 (0% Removal Efficiency)

SWC: 347.71 ppb

DW: LADD: 1.14×10^{-5} mg/kg/d, ADD: 2.85×10^{-5} mg/kg/d, ADR: 1.74×10^{-2} mg/kg/d

Fish: LADD: 1.55×10^{-7} mg/kg/d, ADD: 3.87×10^{-7} mg/kg/d, ADR: 4.33×10^{-4} mg/kg/d

>COC (1,000 ppb): No Exceedance

Use:

10 site, 0 workers, 350 d/yr

Inhalation: Negligible

Dermal: Not Required per SAT

Releases to Water: 6.0×10^{-1} kg/s/d, 1 d/yr

OR Incineration OR Landfill

Releases to Incineration: 2.8×10^{-1} kg/yr

Fate: Releases to Water (0% Removal Efficiency)

SWC: 83.45 ppb

DW: LADD: 3.91×10^{-7} mg/kg/d, ADD: 9.79×10^{-7} mg/kg/d, ADR: 4.19×10^{-3} mg/kg/d

Fish: LADD: 5.30×10^{-9} mg/kg/d, ADD: 1.33×10^{-8} mg/kg/d, ADR: 1.04×10^{-4} mg/kg/d

>COC (1,000 ppb): No Exceedance

PART VII: FOCUS DECISION AND RATIONALE

DISPOSITION: LVE Final Conditional Grant

RATIONALE: L07-0295 was given a final conditional grant based on binding to the production volume of 1,000 kg/yr. Potential risks to human health were address by adequate respiratory protection. The Inhalation Monitoring Criteria for the Pilot Program were met for inhalation exposures from processing. No Inhalation Monitoring is requested. Concerns for potential risks to the environment were low based on low toxicity.

P2REC: This case was nominated for P2 recognition based on its intended use to replace radionuclides tracers used to measure the flow rate in oil-bearing strata and to adjust their pumping rate to achieve desirable flow characteristics. Its use will result in a reduced handling of radioactive materials by both contractors and oil production employees, as well as lower release of radioactive materials into the environment. The Focus participants decided to forward this claim.

PART VIII: CCD DISPOSITION / DD

CCD: